

## Medical Student Distress: Causes, Consequences, and Proposed Solutions

LISELOTTE N. DYRBYE, MD; MATTHEW R. THOMAS, MD; AND TAIT D. SHANAFELT, MD

The goal of medical education is to graduate knowledgeable, skillful, and professional physicians. The medical school curriculum has been developed to accomplish these ambitions; however, some aspects of training may have unintended negative effects on medical students' mental and emotional health that can undermine these values. Studies suggest that mental health worsens after students begin medical school and remains poor throughout training. On a personal level, this distress can contribute to substance abuse, broken relationships, suicide, and attrition from the profession. On a professional level, studies suggest that student distress contributes to cynicism and subsequently may affect students' care of patients, relationship with faculty, and ultimately the culture of the medical profession. In this article, we review the manifestations and causes of student distress, its potential adverse personal and professional consequences, and proposed institutional approaches to decrease student distress.

*Mayo Clin Proc.* 2005;80(12):1613-1622

Medical schools are responsible for ensuring that graduates are knowledgeable, skillful, and professional.<sup>1,2</sup> To achieve these goals, medical schools typically use a curriculum of didactic lectures, modeling, supervised practice, mentoring, and hands-on experience to augment individual study. Unfortunately, some aspects of the training process have unintended negative consequences on students' personal health. Studies suggest that medical students experience a high incidence of personal distress,<sup>3-17</sup> with potential adverse consequences on academic performance,<sup>18-20</sup> competency,<sup>21</sup> professionalism,<sup>22-29</sup> and health.<sup>14,30-34</sup> It is critical for medical educators to understand the prevalence and causes of student distress, potential adverse personal and professional consequences, and institutional factors that can positively and negatively influence student health. In this article, we summarize the manifestations, causes, and consequences of student distress; propose how medical schools can address this problem; and outline areas where additional research is needed.

### METHODS

The intent of this work was to summarize the central themes of medical student distress reported in the literature and to highlight selected studies exploring the prevalence, causes, and consequences of student distress as well as strategies to reduce student distress and promote well-being. Articles were identified by searching MEDLINE and PubMed for English language articles published be-

tween 1966 and 2004 with use of the search terms *medical student AND depression, suicide, stress, burnout, distress, abuse, alcohol drinking, illicit drug usage, street drugs, substance-related disorders, ethics, professionalism, cynicism, cheating, debt, or academic performance*. Additional studies were identified from the reference lists of these articles. Articles were reviewed critically by authors and included as appropriate to provide readers an overview of the research on medical student distress to date, with specific works featured based on the validity of methods used, the novelty of the research question, and the clarity of the findings. As such, this work is intended to be a summary rather than a systematic review that gives readers an understanding of the current literature on medical student distress.

### MANIFESTATIONS OF STUDENT DISTRESS

#### STRESS

Medical students experience substantial stress from the beginning of the training process.<sup>3,5,35-42</sup> Although some degree of stress is a normal part of medical training and can be a motivator for some individuals, not all students find stress constructive.<sup>43</sup> For many individuals, stress arouses feelings of fear, incompetence, uselessness, anger, and guilt and can be associated with both psychological<sup>3,4,10,38,44,45</sup> and physical morbidity.<sup>10,46</sup>

Students use various coping mechanisms to process stress that vary by year in training and source of stress.<sup>47</sup> The specific coping strategies that students use may determine the effect of stress on psychological and physical health<sup>4,10,12,38,46,48</sup> and may determine whether stress has a positive or negative influence.<sup>4,12,47,49</sup> Strategies that center on disengagement, such as problem avoidance, wishful thinking, social withdrawal, and self-criticism, have negative consequences and correlate with depression, anxiety,

From the Department of Internal Medicine and Division of Primary Care Internal Medicine (L.N.D., M.R.T.) and Department of Internal Medicine and Division of Hematology (T.D.S.), Mayo Clinic College of Medicine, Rochester, Minn.

Individual reprints of this article are not available. Address correspondence to Liselotte N. Dyrbye, MD, Division of Primary Care Internal Medicine, 200 First St SW, Rochester, MN 55905 (e-mail: dyrbye.liselotte@mayo.edu).

© 2005 Mayo Foundation for Medical Education and Research

and poor mental health.<sup>4,38,45</sup> In contrast, strategies that involve engagement, such as problem solving, positive reinterpretation, reliance on social support, and expression of emotion, enable students to respond in a manner that leads to adaptation,<sup>4,10,38,46</sup> which can reduce anxiety and depression and their effects on mental<sup>38</sup> and physical health.<sup>46</sup>

#### DEPRESSION

At the start of medical school, medical students have mental health similar to their nonmedical peers.<sup>9,17</sup> Given that the aims of medical training include teaching graduates how to “promote health” and prepare for a career in an intellectually stimulating and socially meaningful profession, it is tempting to speculate that medical school would be a time of personal growth and enhanced health. Unfortunately, the contrary appears to be true, with numerous studies suggesting that students’ mental health worsens during medical school.<sup>3,17</sup>

Surveys in both the United States and abroad identify a high frequency of depression and poor mental health among medical students.<sup>3-10</sup> In a study from the United Kingdom, more than one third of first-year students had poor mental health when measured with the General Health Questionnaire 12, which assesses anxiety and depression.<sup>3</sup> Another study from the United Kingdom of that year students found that the incidence of poor mental health on the General Health Questionnaire 12 doubled during the first year, increasing from 25% to 52%.<sup>4</sup> Other studies have reported similar findings<sup>5</sup> and suggest that this decline in students’ mental health persists throughout the remainder of medical school.<sup>6</sup>

In a 2002 survey of first- and second-year US medical students, 24% of students were depressed according to the Beck Depression Inventory.<sup>7</sup> In a separate study, median Beck Depression Inventory scores increased 3-fold from the time of matriculation to the end of the second year, with 25% of students dysphoric, if not clinically depressed.<sup>8</sup> Two additional studies of US students confirm a peak in depression during the second year of medical school, with gradual improvement during the third and fourth year of training.<sup>9,10</sup>

Despite the high prevalence of mental health–related concerns and ready access to mental health services, depressed medical students are no more likely than the general population to seek treatment for depression.<sup>7,11,50,51</sup> Few students use mental health services,<sup>7,51</sup> instead relying on the support of family and friends during periods of mental illness.<sup>51</sup> Barriers to use of mental health services include lack of time, perception of academic jeopardy, concern regarding confidentiality, the stigma of mental illness, and cost.<sup>7,11,51</sup> Some of these issues appear to be even greater barriers for female and minority students.<sup>11</sup>

#### BURNOUT

Burnout is another measure of poor mental health attributed to work-related stress. This syndrome of emotional exhaustion, depersonalization, and low personal accomplishment culminates in decreased effectiveness at work and is particularly common in individuals in the helping professions (teachers, nurses, social workers, etc).<sup>52</sup> Several studies have reported high rates of burnout in residents<sup>53,54</sup> and practicing physicians,<sup>55-57</sup> leading to speculation that the origin of physician burnout occurs during medical school.<sup>27,58</sup> Despite this hypothesis, the prevalence, inciting factors, and effect of burnout among medical students have not been well studied,<sup>6</sup> particularly among US students.

### POTENTIAL CAUSES OF STUDENT DISTRESS

#### ADJUSTMENT TO THE MEDICAL SCHOOL ENVIRONMENT

The sources of stress for medical students vary by year in training. The first-year medical student is faced with the challenges of being uprooted from family and friends and adapting to a demanding new learning environment. Human cadaver dissection is a well-recognized stress for many students,<sup>3</sup> but other sources of distress, such as a substantially increased scholastic workload<sup>3,49</sup> and concern for academic performance,<sup>41</sup> also characterize this transition. Attempting to master a large volume of information and joining a peer group of equal motivation and intelligence can be intimidating for young adults accustomed to rapid mastery of material and academic distinction. This challenge is amplified for students who struggle academically.<sup>39</sup> High-stakes examinations, such as the part I examination of the National Board of Medical Examiners<sup>3,42</sup> and tests that must be passed before academic advancement, frequently lead to performance anxiety at the end of the preclinical years.<sup>59</sup>

Once in the clinical years of training, students often are separated from their peer-support group and frequently rotate to new work environments at different hospitals. Each rotation requires a unique medical knowledge base and skill set, which tends to highlight students’ deficiencies rather than their progress.<sup>60</sup> An unstructured learning environment, lack of time for recreation, concerns about financial issues, long on-duty assignments, student abuse, and exposure to human suffering can be additional sources of distress during this period.<sup>43,49,60-63</sup>

#### ETHICAL CONFLICTS

Interpersonal interactions between learners (students) and teachers (faculty, residents, and interns) can subtly but profoundly influence students.<sup>64</sup> The vast majority of these interactions are between students and residents and occur in an informal setting when no faculty are present.<sup>65</sup> This

“informal curriculum” conveys powerful messages about professional values, character, and norms. Unfortunately, depression, burnout, and stress are common among student supervisors<sup>65,66</sup> and can lead to modeling of cynicism and unethical behavior that contradict lessons from the formal curriculum.<sup>65,67-70</sup>

In 1 study of third- and fourth-year students, 98% of respondents reported observing physicians refer to patients in a derogatory fashion, and 60% reported witnessing unethical behavior toward a patient.<sup>67</sup> More than two thirds of students experienced guilt about their personal role in these episodes but felt forced to participate to “fit in” and receive a favorable evaluation. Others have made similar observations.<sup>70,71</sup> The “see one, do one, teach one” approach to learning invasive procedures, the demands to write notes about patients not personally examined, and a dehumanized approach to patients (“divide and conquer”) also can present ethical challenges to students who desire to be “team players” and who are influenced strongly by supervising physicians.<sup>65,67,69-73</sup>

#### EXPOSURE TO DEATH AND HUMAN SUFFERING

Most patients receive much of their health care toward the end of life, and medical students in the clinical years are confronted frequently with issues related to death and dying for the first time.<sup>74</sup> Unfortunately, the medical school curriculum often focuses exclusively on disease diagnosis and treatment and pays little attention to education about end-of-life issues and palliative care.<sup>62,74,75</sup> In light of the frequency with which students encounter patients at the end of life and the lack of student training in this area, it is no surprise that students are fearful, anxious, and hesitant to interact with dying patients.<sup>74</sup> Students report feeling awkward, sad, overwhelmed, apprehensive, vulnerable, angry, and anxious in these circumstances,<sup>62,75</sup> which highlights the limitations of medical science and can precipitate thoughts about one’s own death.<sup>62,74,76-78</sup>

Medical students recognize that they typically are inadequately prepared to communicate with dying patients and their families.<sup>63,74</sup> Although issues related to death and dying often are presented during preclinical lectures, clinical training in the skills required to care for patients at the end of life is less common.<sup>62,63</sup> One study reported that although 100% of third-year students had cared for a terminally ill patient, only 41% had been present while an attending physician talked with a dying patient, and only 35% had ever discussed with an attending physician how to care for terminally ill patients.<sup>63</sup>

#### STUDENT ABUSE

The perception of being taken advantage of or abused is common (50%-85%) among medical students.<sup>61,79-81</sup> More

female than male medical students report having been victims of abuse,<sup>80,82-84</sup> but no variation by race has been reported in the few studies published.<sup>82,83</sup> Student abuse occurs most often during the clinical years,<sup>61,85</sup> with faculty, house staff, and nurses the most common abusers.<sup>61,80,83-86</sup>

Although verbal abuse is the most common problem in this category,<sup>61,80,81,85,86</sup> institutional abuse (unfair grades, excessive workload, unnecessary scut work),<sup>61</sup> assignment of inappropriate task (ie, getting food for the team),<sup>86</sup> physical abuse,<sup>61,81,85,86</sup> sexual harassment,<sup>61,79,81,84-86</sup> and racial discrimination<sup>79,81,84</sup> are also serious problems. The effect of abuse on students is serious.<sup>80-84,87</sup> In 1 study of more than 500 medical students, more than 40% reported that they had personally experienced abuse, with many stating that the experience was a major source of stress that affected them for a month or longer.<sup>61</sup>

Regardless of year in training, verbal abuse seriously affects students’ confidence<sup>82-84,87</sup> and negatively affects the learning environment.<sup>81-83</sup> Studies also suggest that verbal abuse influences students’ specialty choice,<sup>82,83</sup> adversely affects their care of patients,<sup>83</sup> decreases institutional loyalty,<sup>83</sup> and erodes mental health.<sup>80,81,83</sup> Anxiety, depression, hostility, low self-esteem, and use of alcohol to “escape”<sup>80</sup> are more common among students who perceive abuse.<sup>80,83</sup> Despite this profound effect, less than one third of students report abuse to faculty or medical school administrators,<sup>79,81,82</sup> often due to fear of reprisal or concern of potential repercussions on performance evaluations.<sup>79,82</sup>

#### PERSONAL LIFE EVENTS

Although sources of stress related to the training experience have been the focus of most research on student distress, students also experience numerous personal life stressors common to individuals their age. In a study of more than 1000 medical students, many reported experiencing the death of a family member (15%), personal illness or injury (25%), or change of health in a relative (42%) within the past year.<sup>88</sup> Although these life events would be expected to adversely affect students’ quality of life (QOL) and professional development, their effect has not been well studied. Other personal life events, such as marriage, appear to protect against distress. In the 1995 Association of American Medical Colleges graduation questionnaire, 30% of graduating medical students were married (a lower prevalence than reported in the age-matched general population),<sup>89</sup> and another 14% were engaged or partnered.<sup>90</sup> The lower stress found among married students relative to their single counterparts<sup>91</sup> has been attributed to emotional support provided by the spouse.<sup>91,92</sup>

Although marriage is relatively common among medical students, smaller numbers of students (10%) have children by graduation,<sup>90</sup> and little is known about the mental

health consequences of pregnancy or childrearing during medical school. Although childbirth and childrearing typically are considered positive life events, children add a level of complexity to students' lives,<sup>93,94</sup> with a mental health effect that may be gender-specific.<sup>9</sup> In 1 study of second-year medical students, female students were more likely to be depressed if they had children, whereas no such relationship was observed among their male colleagues.<sup>9</sup>

#### EDUCATIONAL DEBT

Student debt has increased out of proportion to the escalation of tuition fees in recent years.<sup>84</sup> The mean educational debt for the class of 2004 was \$115,218 (up 5% from the previous year).<sup>79</sup> Student debt among Canadian graduates also is increasing, leading some graduates to leave home to pursue more lucrative careers in the United States.<sup>95</sup> Although the effect of debt on medical students' mental health has not been studied directly, financial concerns are a common source of student stress<sup>49</sup> and appear to influence specialty choice.<sup>96</sup>

### CONSEQUENCES OF STUDENT DISTRESS

#### IMPAIRED ACADEMIC PERFORMANCE

Stress and performance are related intimately. Performance measures such as grades and clinical evaluations can be sources of stress and anxiety.<sup>39</sup> Although these states of distress may reciprocally affect grades,<sup>20</sup> the degree of their influence may be subtle and depend on personality.<sup>97-100</sup> In 1 study, both grades in the preclinical years and clerkship examination scores could be predicted as well by psychosocial characteristics (anxiety, depression, loneliness, neuroticism, self-esteem, and stressful life events) as by Medical College Admissions Test scores.<sup>20</sup> Even more surprising, psychosocial characteristics were better predictors of clinical competency than the admissions test scores.<sup>20</sup> Specific stressors, such as interpersonal conflicts with team members, also appear to influence clerkship grades<sup>18,19</sup> and National Board of Medical Examiners part II examination scores.<sup>18</sup> Most<sup>101-103</sup> but not all<sup>39</sup> studies also suggest that anxiety correlates with poor performance, but cause vs effect is difficult to determine.<sup>103</sup> Although theorized to have a negative influence, the effect of depression and burnout on academic performance in medical students has not been well studied.<sup>104</sup>

#### CYNICISM

Although "an interest in helping people" is one of the most common reasons that college graduates cite for choosing a career in medicine,<sup>105</sup> this idealism often gives way to cynicism during medical school.<sup>24-29</sup> The medical school training process actually is characterized by a decline in

empathy and humanitarianism<sup>22,24</sup>—2 traits that medical educators strive to promote. This decline in compassion, initially recognized in the 1950s,<sup>28</sup> begins during the preclinical years and progresses during clinical rotations.<sup>22</sup> Such negative attitudes may develop in response to students' environment and experiences. Although in the short term, attitudes such as cynicism may serve as a buffer against anxiety, fear of failure, and exposure to human suffering,<sup>27,73,106</sup> they ultimately erode professionalism. Empathy has been shown to correlate with physician competency,<sup>21</sup> and the erosion of this characteristic throughout the course of training is particularly concerning.<sup>27,29,67,107-109</sup> Cynicism and loss of compassion also appear to affect specialty choice<sup>110-113</sup> and can translate into an unwillingness to care for chronically ill, elderly, and terminal patients.<sup>112,114</sup> Increasingly, students also appear to be basing their specialty choice on lifestyle considerations rather than on humanitarian ideals,<sup>96</sup> reflecting both a change in priorities and the desire to limit personal distress.<sup>113</sup>

#### ACADEMIC DISHONESTY

A lack of integrity among some medical school applicants, medical students, residents, and physicians has been well described.<sup>115-118</sup> Nearly one quarter of students admit to cheating,<sup>117</sup> and more than two thirds report witnessing cheating by colleagues.<sup>116</sup> Students cite illness, workload, and perception of the material taught as "trivial" as reasons they cheat.<sup>117</sup> Dishonesty in patient care activities, such as recording tasks not performed, reporting findings elicited by others, and lying about having ordered a test, often are motivated by fear and a desire to appear knowledgeable.<sup>117</sup>

The perception of what defines academic integrity also differs by year in school. In 1 cross-sectional study, first-year students were more likely than more senior students to correctly identify case scenarios describing academic dishonesty as being unacceptable.<sup>119</sup> Students in later years of training were both less likely to consider the behaviors wrong and more likely to report that they had or would engage in the behaviors described.<sup>119</sup>

#### SUBSTANCE ABUSE

Although the overall pattern of alcohol consumption among medical students is similar to that of age-related peers,<sup>120,121</sup> problematic alcohol consumption is common among medical students.<sup>14,30-34</sup> Up to 20% of first-year medical students admit to excessive alcohol intake<sup>14,30,32</sup> and report anxiety, stress, examination and work pressures, and tension to be among the common reasons for alcohol consumption.<sup>34,122,123</sup> At one Midwest medical school, students' mean score on a validated assessment for alcohol dependence (Alcohol Use Disorders Identification Test)



doubled during the first semester ( $P < .001$ ), with 20% of students scoring above the cutoff for problematic drinking.<sup>14</sup> Other studies suggest that alcohol use remains high throughout medical training.<sup>14,32,34,123</sup> In a study of graduating students from 8 US medical schools, 29% of students reported that their alcohol consumption increased during medical school, with more than 20% reporting at least 1 episode of binge drinking (5 or more drinks in 1 sitting) in the past 30 days.<sup>33</sup>

Illicit drug use among medical students is comparable to or less than that of age-related peers.<sup>31,122</sup> Despite this fact, 3% to 10% of medical students report using illicit substances.<sup>31</sup> In a survey of 2046 senior students at 23 medical schools, the reported incidence of marijuana (10%), cocaine (2.8%), tranquilizer (2.3%), heroin/opiate (1.1%), psychedelic (0.7%), amphetamine (0.3%), and barbiturate (0.2%) use in the last 30 days was concerning.<sup>122</sup> Most,<sup>122,124</sup> but not all,<sup>34</sup> studies suggest that students who report use of illicit drugs started this practice before medical school.

#### SUICIDE

Suicide is the third-leading cause of death among 20- to 30-year-olds in the United States.<sup>125</sup> The annual suicide rate for male medical students between 1974 and 1981 was comparable to similarly aged men in the general population.<sup>126</sup> Although the suicide rate for female students during this period equaled that of their male colleagues, it was 3 to 4 times higher than age-matched women in the general population.<sup>126</sup> In a follow-up study of 101 US medical schools, 15 students were reported to have committed suicide between August 1989 and May 1994.<sup>127</sup> All but 1 of these suicides were committed by men.<sup>127</sup> In this study, suicide ranked higher as a cause of death in medical students than in similarly aged Americans.<sup>127</sup>

Although the prevalence of suicidal ideation and planning among medical students has not been well studied,<sup>15</sup> an estimated 8 to 25 attempted suicides occur for each suicide death, reflecting the concern that completed suicides represent only a fraction of the extreme distress manifested by suicidal ideation, planning, and attempts among medical students.<sup>125,128,129</sup> The risk of student suicide appears higher in the clinical years. Among senior Norwegian medical students, 14% reported having suicidal thoughts within the past year, and 6% had planned to commit suicide during medical school. Such suicidal thoughts persist into postgraduate training<sup>15</sup> and practice.<sup>130</sup>

Depression, personal life events, and personality traits may influence the transition from suicidal ideation to planning,<sup>131</sup> but the factors that prompt medical students to act on their plan have not been studied. Among physicians, suicide attempts are more likely among those who are

single, female, depressed, have other psychiatric illness, or struggle with drug or alcohol dependency.<sup>128,132</sup>

### REDUCING STUDENT DISTRESS: IDEAS FOR MEDICAL SCHOOLS AND MEDICAL EDUCATORS

Understanding the causes and consequences of student distress is important (Figure 1), but medical schools need to go beyond identifying distress and strive to promote well-being for all students. Well-being is distinct from the mere absence of distress and includes achieving a high QOL in multiple domains (physical health, mental health, emotional health, spiritual health, etc).<sup>133,134</sup> Promoting and nurturing well-being during medical school and equipping graduates with the skills necessary to recognize personal distress, to determine when they need to seek assistance, and to develop strategies to promote their own well-being is essential to promoting professionalism and laying the foundation for resilience through the course of a career.

#### CREATING A NURTURING LEARNING ENVIRONMENT

The Association of American Medical Colleges urges medical schools to establish relationships between faculty members and students to promote a positive learning environment.<sup>135</sup> Examples and characteristics of positive faculty-student mentoring programs have been described,<sup>136-138</sup> with studies suggesting that these programs have a positive effect on student well-being.<sup>139,140</sup> Although relationships with faculty undoubtedly assist students, student-led support programs may provide an even stronger source of support and promote positive strategies for coping with stress.<sup>141</sup> Senior medical students may more easily relate to challenges faced by junior students, and “buddy programs” designed to promote mentorship of junior students by senior students appear to lower student stress.<sup>142</sup> Fostering relationships between classes and with faculty through institution-sponsored social events also can reduce stress and help prevent burnout.<sup>16,53,143-145</sup>

The evaluation system used to assess student performance also can have a powerful effect on the learning environment. The A-F grading scheme, used to classify performance, often creates a competitive environment that promotes anxiety and peer competition rather than collaborative learning. Researchers at the University of Michigan evaluated the effect of changing to a pass-fail grading system on student performance and satisfaction.<sup>146</sup> Compared with previous classes, students’ performance on tests in basic science courses were unchanged, suggesting students’ motivation for subject mastery was not affected by the change in the evaluation system. In contrast, students’ satisfaction with the evaluation system and learning envi-




FIGURE 1. Proposed model of causes and consequences of student distress.

ronment improved with the pass-fail approach. Although some authors have reported that residency program directors prefer candidates evaluated with use of traditional grades,<sup>147,148</sup> others have reported that the pass-fail grading system does not influence students' likelihood of matching with a highly ranked postgraduate training program.<sup>149</sup>

Aligning the "informal curriculum" with the ideals of compassion and professionalism is also essential to create a positive learning environment.<sup>71</sup> Thus, fostering a nurturing environment for students in part may depend on promoting the well-being of residents and faculty. Unfortunately, burnout and cynicism are common among residents and practicing physicians and can adversely affect professional modeling.<sup>52,55-57</sup> Faculty development programs need to both address staff satisfaction and confront problematic behavior such as disrespect, hostility, and rudeness, which are often ignored.<sup>150</sup>

Finally, autonomy is a central component of physician job satisfaction and is likely important to students.<sup>57,151</sup> Allowing students to contribute to curriculum development can benefit both students and administrators and give students a sense of ownership in their educational experience. Students bring unique perspectives to curriculum committees including insight regarding redundancies in the curriculum, feedback on effective and ineffective teaching methods, and ideas about areas for new curriculum devel-

opment (alternative medicine, end-of-life care, ethics, genomics, etc). Because students rotate among various hospitals in the training system, they also can provide insight regarding variations in the care of patients, workload, culture, and teaching style among hospitals and identify the most effective experiences.<sup>69,73</sup>

**IDENTIFYING AND ASSISTING STRUGGLING STUDENTS**

Poor undergraduate scholastic performance,<sup>38,39</sup> poor academic performance early in medical school,<sup>38,45</sup> mental health problems,<sup>38,45,46</sup> and avoidant coping mechanisms<sup>38,46</sup> place students at risk of distress both during and after medical school.<sup>38,46</sup> Simply making students aware of their mental health "profile" does not appear to reduce distress,<sup>14</sup> and once struggling students are identified, they need individualized support. Deans must not only make students aware of the available resources, but also address barriers to care. Because of the stigma of mental illness, many medical students are not comfortable seeking care for mental health problems from their own institution and prefer to receive off-site care.<sup>11,51</sup> Creating an ombudsman program, offering career counseling, and providing students off-campus confidential resources covered by the student health insurance plan are critical. Descriptions of successful mental health programs for medical students may serve as models.<sup>145,152-157</sup>

### TEACHING SKILLS FOR STRESS MANAGEMENT AND PROMOTING SELF-AWARENESS

Teaching students to use adaptive coping mechanisms,<sup>48</sup> such as acceptance, planning, positive reinterpretation, and self-distraction, can reduce psychological morbidity.<sup>4,10,38</sup> Stress-management programs that inform students about the effects of stress on physiological and psychological functioning and teach students how to plan, prioritize, identify sources of stress, and cope with stress<sup>144,145,158,159</sup> reduce tension and anxiety and simultaneously increase awareness and use of positive coping strategies.<sup>144,158-160</sup> Peer discussion groups can help students process conflict, nurture self-awareness, and promote empathy.<sup>161,162</sup> Such groups also provide opportunities for students to express, analyze, and share feelings, which decreases the likelihood of burnout.<sup>52</sup> Such shared reflection helps students realize that their struggles are not unique<sup>162</sup> and provides insight into how colleagues solve similar problems.

### HELPING STUDENTS PROMOTE PERSONAL HEALTH

Encouraging students to promote personal health with regular physical activity and adequate sleep is also valuable.<sup>16,48,163</sup> Requiring students to establish a primary care physician at matriculation facilitates delivery of age-appropriate preventive services and establishes a care provider with whom students can discuss distress, mental health problems, and substance use. Students should also be provided appropriate time off during holidays and between rotations to allow them to decompress from the rigors of training. Explicit promotion of work-life balance, including role-modeling by medical school faculty, also may be of great benefit. Presentations by faculty regarding how they have personally handled challenges in work-life balance are an excellent way to expose students to this concept and stimulate discussion.

### AREAS OF NEEDED RESEARCH

Although much is known about student distress, little is known about student well-being, how it can be fostered, and its potential to enhance learning and professional development. Validated tools are available to measure various domains of QOL and should be used for such studies.<sup>164-171</sup> Research is needed to identify personal and program factors that enhance students' QOL (physical health, emotional health, spiritual health, etc), assess the efficacy of support systems to assist struggling students, and develop curriculum to promote well-being. Prospective longitudinal studies exploring changes in student stress, burnout, and QOL from the time of matriculation through the completion of residency training would be

enlightening. The influence of faculty distress and QOL on student QOL and professional development also needs to be explored.

### CONCLUSIONS

Medical school is a stressful period of physician training. Many medical students experience substantial distress, which contributes to poor academic performance, academic dishonesty, cynicism, and substance abuse. Medical educators need to be aware of the manifestations, causes, and consequences of student distress, and medical schools need to develop and evaluate programs to support struggling students and promote student well-being. Additional research is needed to identify personal and program factors that promote well-being and explore its potential to enhance competency. In the long run, efforts to promote students' well-being will benefit patients, the public, and the profession, in addition to the individual.

*We are grateful to Dr Jarrett W. Richardson for his critical review of the submitted manuscript.*

### REFERENCES

1. Liaison Committee on Medical Education. Functions and Structure of a Medical School: Standards for Accreditation of Medical Education Programs Leading to the M.D. Degree. March 2003. Available at: [www.lcme.org/functions2003march.pdf](http://www.lcme.org/functions2003march.pdf). Accessibility verified October 24, 2005.
2. Association of American Medical Colleges. Report I. Learning Objectives for Medical Student Education: Guidelines for Medical Schools. Medical Schools Objectives Project. January 1998. Available at: [www.aamc.org/meded/msop/msop1.pdf](http://www.aamc.org/meded/msop/msop1.pdf). Accessibility verified October 24, 2005.
3. Guthrie EA, Black D, Shaw CM, Hamilton J, Creed FH, Tomenson B. Embarking upon a medical career: psychological morbidity in first year medical students. *Med Educ*. 1995;29:337-341.
4. Moffat KJ, McConnachie A, Ross S, Morrison JM. First year medical student stress and coping in a problem-based learning medical curriculum. *Med Educ*. 2004;38:482-491.
5. Aktekin M, Karaman T, Senol YY, Erdem S, Erengin H, Akaydin M. Anxiety, depression and stressful life events among medical students: a prospective study in Antalya, Turkey. *Med Educ*. 2001;35:12-17.
6. Guthrie E, Black D, Bagalkote H, Shaw C, Campbell M, Creed F. Psychological stress and burnout in medical students: a five-year prospective longitudinal study. *J R Soc Med*. 1998;91:237-243.
7. Givens JL, Tjia J. Depressed medical students' use of mental health services and barriers to use. *Acad Med*. 2002;77:918-921.
8. Clark DC, Zeldow PB. Vicissitudes of depressed mood during four years of medical school. *JAMA*. 1988;260:2521-2528.
9. Rosal MC, Ockene IS, Ockene JK, Barrett SV, Ma Y, Hebert JR. A longitudinal study of students' depression at one medical school. *Acad Med*. 1997;72:542-546.
10. Mosley TH Jr, Perrin SG, Neral SM, Dubbert PM, Grothues CA, Pinto BM. Stress, coping, and well-being among third-year medical students. *Acad Med*. 1994;69:765-767.
11. Roberts LW, Warner TD, Lyketsos C, Frank E, Ganzini L, Carter D. Collaborative Research Group on Medical Student Health. Perceptions of academic vulnerability associated with personal illness: a study of 1,027 students at nine medical schools. *Compr Psychiatry*. 2001;42:1-15.
12. Vitaliano PP, Maiuro RD, Russo J, Mitchell ES, Carr JE, Van Citters RL. A biopsychosocial model of medical student distress. *J Behav Med*. 1988;11:311-331.
13. Raj SR, Simpson CS, Hopman WM, Singer MA. Health-related quality of life among final-year medical students. *CMAJ*. 2000;162:509-510.

14. Ball S, Bax A. Self-care in medical education: effectiveness of health-habits interventions for first-year medical students. *Acad Med.* 2002;77:911-917.
15. Tyssen R, Vaglum P, Gronvold NT, Ekeberg O. Suicidal ideation among medical students and young physicians: a nationwide and prospective study of prevalence and predictors. *J Affect Disord.* 2001;64:69-79.
16. Parkerson GR Jr, Broadhead WE, Tse CK. The health status and life satisfaction of first-year medical students. *Acad Med.* 1990;65:586-588.
17. Carson AJ, Dias S, Johnston A, et al. Mental health in medical students: a case control study using the 60 item General Health Questionnaire. *Scott Med J.* 2000;45:115-116.
18. Spiegel DA, Smolen RC, Jonas CK. An examination of the relationships among interpersonal stress, morale and academic performance in male and female medical students. *Soc Sci Med.* 1986;23:1157-1161.
19. Spiegel DA, Smolen RC, Hopfensperger KA. Medical student stress and clerkship performance. *J Med Educ.* 1986;61:929-931.
20. Hojat M, Robeson M, Damjanov I, Veloski JJ, Glaser K, Gonnella JS. Students' psychosocial characteristics as predictors of academic performance in medical school. *Acad Med.* 1993;68:635-637.
21. Hojat M, Gonnella JS, Mangione S, et al. Empathy in medical students as related to academic performance, clinical competence and gender. *Med Educ.* 2002;36:522-527.
22. Woloschuk W, Harasym PH, Temple W. Attitude change during medical school: a cohort study. *Med Educ.* 2004;38:522-534.
23. Hojat M, Mangione S, Nasca T, et al. An empirical study of decline in empathy in medical school. *Med Educ.* 2004;38:934-941.
24. Crandall SJ, Volk RJ, Loemker V. Medical students' attitudes toward providing care for the underserved: are we training socially responsible physicians? *JAMA.* 1993;269:2519-2523.
25. Rezler AG, Ten Haken JT. Affect and research in medical education. *Med Educ.* 1984;18:331-338.
26. Eron L. The effect of medical education on attitudes: a follow-up study. *J Med Educ.* 1958;33:25-33.
27. Wolf TM, Balson PM, Faucett JM, Randall HM. A retrospective study of attitude change during medical education. *Med Educ.* 1989;23:19-23.
28. Eron L. Effect of medical education on medical students' attitudes. *J Med Educ.* 1955;30:559-566.
29. Rezler AG, Flaherty JA. *The Interpersonal Dimension in Medical Education.* New York, NY: Springer; 1985:147-182.
30. Gutgesell M, Reeve R, Parsons B, Morse R. Exercise and alcohol consumption among medical students. *Acad Med.* 1999;74:750-751.
31. Croen LG, Woensner M, Herman M, Reichgott M. A longitudinal study of substance use and abuse in a single class of medical students. *Acad Med.* 1997;72:376-381.
32. Clark D, Eckenfels EJ, Daugherty SR, Fawcett J. Alcohol-use patterns through medical school: a longitudinal study of one class. *JAMA.* 1987;257:2921-2926.
33. Mangus R, Hawkins C, Miller M. Tobacco and alcohol use among 1996 medical school graduates. *JAMA.* 1998;280:1192-1193.
34. Newbury-Birch D, Walshaw D, Kamali F. Drink and drugs: from medical students to doctors. *Drug Alcohol Depend.* 2001;64:265-270.
35. Daly MG, Willcock SM. Examining stress and responses to stress in medical students and new medical graduates. *Med J Aust.* 2002;177(suppl):S14-S15.
36. Kiessling C, Schubert B, Scheffner D, Burger W. First year medical students' perceptions of stress and support: a comparison between reformed and traditional track curricula. *Med Educ.* 2004;38:504-509.
37. Toews JA, Lockyer JM, Dobson DJ, et al. Analysis of stress levels among medical students, residents, and graduate students at four Canadian schools of medicine. *Acad Med.* 1997;72:997-1002.
38. Stewart SM, Betson C, Lam TH, Marshall IB, Lee PW, Wong CM. Predicting stress in first year medical students: a longitudinal study. *Med Educ.* 1997;31:163-168.
39. Stewart SM, Lam TH, Betson CL, Wong CM, Wong AM. A prospective analysis of stress and academic performance in the first two years of medical school. *Med Educ.* 1999;33:243-250.
40. Firth J. Levels and sources of stress in medical students. *Br Med J (Clin Res Ed).* 1986;292:1177-1180.
41. Supe AN. A study of stress in medical students at Seth G.S. Medical College. *J Postgrad Med.* 1998;44:1-6.
42. Rosenthal TL, Rosenthal RH, Edwards NB. Students' self-ratings of stress in medical school: a replication across 20 months. *Behav Res Ther.* 1990;28:171-173.
43. Linn BS, Zeppa R. Stress in junior medical students: relationship to personality and performance. *J Med Educ.* 1984;59:7-12.
44. Bramness JG, Fixdal TC, Vaglum P. Effect of medical school stress on the mental health of medical students in early and late clinical curriculum. *Acta Psychiatr Scand.* 1991;84:340-345.
45. Tyssen R, Vaglum P, Gronvold NT, Ekeberg O. Factors in medical school that predict postgraduate mental health problems in need of treatment: a nationwide and longitudinal study. *Med Educ.* 2001;35:110-120.
46. Park CL, Adler NE. Coping style as a predictor of health and well-being across the first year of medical school. *Health Psychol.* 2003;22:627-631.
47. Stern M, Norman S, Komm C. Medical students' differential use of coping strategies as a function of stressor type, year of training, and gender. *Behav Med.* 1993;18:173-180.
48. Wolf TM. Stress, coping and health: enhancing well-being during medical school. *Med Educ.* 1994;28:8-17.
49. Wolf TM, Faucett JM, Randall HM, Balson PM. Graduating medical students' ratings of stresses, pleasures, and coping strategies. *J Med Educ.* 1988;63:636-642.
50. Hirschfeld RM, Keller MB, Panico S, et al. The National Depressive and Manic-Depressive Association consensus statement on the undertreatment of depression. *JAMA.* 1997;277:333-340.
51. Chew-Graham CA, Rogers A, Yassin N. 'I wouldn't want it on my CV or their records': medical students' experiences of help-seeking for mental health problems. *Med Educ.* 2003;37:873-880.
52. Maslach C. Burned out. *Hum Behav.* 1976;5:16-22.
53. Lemkau JP, Purdy RR, Rafferty JP, Rudisill JR. Correlates of burnout among family practice residents. *J Med Educ.* 1988;63:682-691.
54. Thomas N. Resident burnout. *JAMA.* 2004;292:2880-2889.
55. Visser MR, Smets EM, Oort FJ, De Haes HC. Stress, satisfaction and burnout among Dutch medical specialists. *CMAJ.* 2003;168:271-275.
56. Deckard GJ, Hicks LL, Hamory BH. The occurrence and distribution of burnout among infectious diseases physicians. *J Infect Dis.* 1992;165:224-228.
57. Linzer M, Visser MR, Oort FJ, Smets EM, McMurray JE, de Haes HC, Society of General Internal Medicine (SGIM) Career Satisfaction Study Group (CSSG). Predicting and preventing physician burnout: results from the United States and the Netherlands. *Am J Med.* 2001;111:170-175.
58. Krakowski A. Stress and the practice of medicine: the myth and the reality. *J Psychosom Res.* 1982;26:91-98.
59. Kidson M, Hornblow A. Examination anxiety in medical students: experiences with the visual analogue scale for anxiety. *Med Educ.* 1982;16:247-250.
60. Colford JM Jr, McPhee SJ. The ravelled sleeve of care: managing the stresses of residency training. *JAMA.* 1989;261:889-893.
61. Silver HK, Glick AD. Medical student abuse: incidence, severity, and significance. *JAMA.* 1990;263:527-532.
62. Wear DP. "Face-to-face with it": medical students' narratives about their end-of-life education. *Acad Med.* 2002;77:271-277.
63. Rappaport W, Witzke D. Education about death and dying during the clinical years of medical school. *Surgery.* 1993;113:163-165.
64. Hafferty FW. Beyond curriculum reform: confronting medicine's hidden curriculum. *Acad Med.* 1998;73:403-407.
65. Stern DT. In search of the informal curriculum: when and where professional values are taught. *Acad Med.* 1998;73(10, suppl):S28-S30.
66. Ramirez AJ, Graham J, Richards MA, Cull A, Gregory WM. Mental health of hospital consultants: the effects of stress and satisfaction at work. *Lancet.* 1996;347:724-728.
67. Feudtner C, Christakis DA, Christakis NA. Do clinical clerks suffer ethical erosion? students' perceptions of their ethical environment and personal development. *Acad Med.* 1994;69:670-679.
68. Madsen T. Resident burnout [letter]. *Ann Intern Med.* 2002;137:698-700.
69. Hafferty FW, Franks R. The hidden curriculum, ethics teaching, and the structure of medical education. *Acad Med.* 1994;69:861-871.
70. Christakis DA, Feudtner C. Ethics in a short white coat: the ethical dilemmas that medical students confront. *Acad Med.* 1993;68:249-254.
71. Hundert EM, Hafferty F, Christakis D. Characteristics of the informal curriculum and trainees' ethical choices. *Acad Med.* 1996;71:624-642.
72. Reynolds PP. Reaffirming professionalism through the education community. *Ann Intern Med.* 1994;120:609-614.
73. Rezler AG. Attitude changes during medical school: a review of the literature. *J Med Educ.* 1974;49:1023-1030.
74. Binienda J, Schwartz K, Gaspar D. Training in end-of-life care through interaction with dying patients. *Acad Med.* 2001;76:517.



75. MacLeod R, Parkin C, Pullon S, Robertson G. Early clinical exposure to people who are dying: learning to care at the end of life. *Med Educ.* 2003;37:51-58.
76. Rucker LMD, Shapiro JP. Becoming a physician: students' creative projects in a third-year IM clerkship. *Acad Med.* 2003;78:391-397.
77. Maxwell TL, Passow ES, Plumb J, Sifri RD. Experience with hospice: reflections from third-year medical students. *J Palliat Med.* 2002;5:721-727.
78. Howells K, Gould M, Field D. Fear of death and dying in medical students: effects of clinical experience. *Med Educ.* 1986;20:502-506.
79. Association of American Medical Colleges. 2004 Medical School Graduation Questionnaire: All Schools Report. Available at: [www.aamc.org/data/gq/allschoolsreports/2004.pdf](http://www.aamc.org/data/gq/allschoolsreports/2004.pdf). Accessibility verified October 24, 2005.
80. Richman JA, Flaherty JA, Rospenda KM, Christensen ML. Mental health consequences and correlates of reported medical student abuse. *JAMA.* 1992;267:692-694.
81. Sheehan H, Sheehan D, White K, Leibowitz A, Baldwin DC Jr. A pilot study of medical student 'abuse': student perceptions of mistreatment and misconduct in medical school. *JAMA.* 1990;263:533-537.
82. Elnicki DM, Curry RH, Fagan M, et al. Medical students' perspectives on and responses to abuse during the internal medicine clerkship. *Teach Learn Med.* 2002;14:92-97.
83. Lubitz RM, Nguyen DD. Medical student abuse during third-year clerkships. *JAMA.* 1996;275:414-416.
84. Kassebaum DG, Cutler ER. On the culture of student abuse in medical school. *Acad Med.* 1998;73:1149-1158.
85. Leenthal A, Kaiserman I, Lernau O. Student abuse in medical school: a comparison of students' and faculty's perceptions. *Isr J Med Sci.* 1996;32:229-238.
86. Elnicki DM, Linger B, Asch E, et al. Patterns of medical student abuse during the internal medicine clerkship: perspectives of students at 11 medical schools. *Acad Med.* 1999;74(10, suppl):S99-S101.
87. Schuchert MK. The relationship between verbal abuse of medical students and their confidence in their clinical abilities. *Acad Med.* 1998;73:907-909.
88. Hojat M, Glaser K, Xu G, Veloski JJ, Christian EB. Gender comparisons of medical students' psychosocial profiles. *Med Educ.* 1999;33:342-349.
89. US Census Bureau. Survey of Income and Program Participation (SIPP): Table 2. Marital Event Transition Rates, by Sex and Age: 2001. Available at: [www.census.gov/population/socdemo/marital-hist/p70-97/tab02.pdf](http://www.census.gov/population/socdemo/marital-hist/p70-97/tab02.pdf). Accessibility verified October 24, 2005.
90. Association of American Medical Colleges. Medical School Graduation Questionnaire: 1995 Summary Report for All Schools. Available at: [www.aamc.org/data/gq/allschoolsreports/1995.pdf](http://www.aamc.org/data/gq/allschoolsreports/1995.pdf). Accessibility verified October 24, 2005.
91. Coombs RH. The effect of marital status on stress in medical school. *Am J Psychiatry.* 1982;139:1490-1493.
92. Katz J, Monnier J, Libet J, Shaw D, Beach S. Individual and crossover effect of stress on adjustment in medical student marriages. *J Marital Fam Ther.* 2000;26:341-351.
93. Lazo R. Family matters. *JAMA.* 1991;265:119.
94. Law JK. Starting a family in medical school. *JAMA.* 1997;277:767.
95. Sullivan P. Mortgage-sized debt the new normal for medical students [published correction appears in *CMAJ.* 2003;169:1147]. *CMAJ.* 2003;169:457-458.
96. Dorsey ER, Jarjoura D, Rutecki GW. Influence of controllable lifestyle on recent trends in specialty choice by US medical students. *JAMA.* 2003;290:1173-1178.
97. Hobfoll SE, Anson O, Antonovsky A. Personality factors as predictors of medical students performance. *Med Educ.* 1982;16:251-258.
98. Ferguson E, James D, O'Hehir F, Sanders A, McManus IC. Pilot study of the roles of personality, references, and personal statements in relation to performance over the five years of a medical degree. *BMJ.* 2003;326:429-432.
99. Lievens F, Coetsier P, De Fruyt F, De Maeseneer J. Medical students' personality characteristics and academic performance: a five-factor model perspective. *Med Educ.* 2002;36:1050-1056.
100. Lipton A, Huxham GJ, Hamilton D. Influence of personality on achievement of medical students. *Br J Med Educ.* 1975;9:215-222.
101. Grover PL, Smith DU. Academic anxiety, locus of control, and achievement in medical school. *J Med Educ.* 1981;56:727-736.
102. Van Der Ploeg H. Relationship of state-trait anxiety to academic performance in Dutch medical students. *Psychol Rep.* 1979;45:223-227.
103. Tooth D, Tonge K, McManus IC. Anxiety and study methods in pre-clinical students: causal relation to examination performance. *Med Educ.* 1989;23:416-421.
104. Clark DC, Daugherty SR, Zeldow PB, Gotterer GS, Hedeker D. The relationship between academic performance and severity of depressed mood during medical school. *Compr Psychiatry.* 1988;29:409-420.
105. Epstein RM, Hundert EM. Defining and assessing professional competence. *JAMA.* 2002;287:226-235.
106. McManus IC, Winder BC, Gordon D. The causal links between stress and burnout in a longitudinal study of UK doctors. *Lancet.* 2002;359:2089-2090.
107. American Board of Internal Medicine. *Project Professionalism*. Philadelphia, Pa: American Board of Internal Medicine; 1995.
108. Hojat M, Mangione S, Gonnella JS, Nasca T, Veloski JJ, Kane G. Empathy in medical education and patient care. *Acad Med.* 2001;76:669.
109. Patenaude J, Niyonsenga T, Fafard D. Changes in students' moral development during medical school: a cohort study. *CMAJ.* 2003;168:840-844.
110. Gill D, Palmer C, Mulder R, Wilkinson T. Medical student career intentions at the Christchurch School of Medicine: the New Zealand Wellbeing, Intentions, Debt and Experiences (WIDE) survey of medical students pilot study: results part II. *N Z Med J.* 2001;114:465-467.
111. Newton BW, Savidge MA, Barber L, et al. Differences in medical students' empathy. *Acad Med.* 2000;75:1215.
112. Davis BE, Nelson DB, Sahler OJ, McCurdy FA, Goldberg R, Greenberg LW. Do clerkship experiences affect medical students' attitudes toward chronically ill patients? *Acad Med.* 2001;76:815-820.
113. Merrill J, Lorimor R, Thornby J, Woods A. Caring for terminally ill persons: comparative analysis of attitudes (thanatophobia) of practicing physicians, student nurses, and medical students. *Psychol Rep.* 1998;83:123-128.
114. Griffith CH, Wilson JF. The loss of idealism throughout internship. *Eval Health Prof.* 2003;26:415-426.
115. Petersdorf RG. A matter of integrity. *Acad Med.* 1989;64:119-123.
116. DeWitt C, Baldwin DC Jr, Daugherty SR, Beverley D, Rowley BD, Schwarz M. Cheating in medical school: a survey of second year students at 31 schools. *Acad Med.* 1996;71:267-273.
117. Dans PE. Self-reported cheating by students at one medical school. *Acad Med.* 1996;71(1, suppl):S70-S72.
118. Anderson RE, Obenshain SS. Cheating by students: findings, reflections, and remedies. *Acad Med.* 1994;69:323-332.
119. Rennie SC, Rudland JR. Differences in medical students' attitudes to academic misconduct and reported behavior across the years—a questionnaire study. *J Med Ethics.* 2003;29:97-102.
120. Wechsler H, Dowdall GW, Maenner G, Gledhill-Hoyt J, Lee H. Changes in binge drinking and related problems among American college students between 1993 and 1997: results of the Harvard School of Public Health College Alcohol Study. *J Am Coll Health.* 1998;47:57-68.
121. National Institute on Alcohol Abuse and Alcoholism. Quick Facts—2004. Available at: [www.niaaa.nih.gov/](http://www.niaaa.nih.gov/). Accessibility verified October 25, 2005.
122. Baldwin DC Jr, Hughes PH, Conard SE, Storr CL, Sheehan DV. Substance use among senior medical students: a survey of 23 medical schools. *JAMA.* 1991;265:2074-2078.
123. Tyssen R, Vaglum P, Aasland OG, Gronvold NT, Ekeberg O. Use of alcohol to cope with tension, and its relation to gender, years in medical school and hazardous drinking: a study of two nation-wide Norwegian samples of medical students. *Addiction.* 1998;93:1341-1349.
124. Schwartz R, Lewis D, Hoffmann N, Kyriazi N. Cocaine and marijuana use by medical students before and during medical school. *Arch Intern Med.* 1990;150:883-886.
125. National Institute on Mental Health. Suicide Facts and Statistics—2004. Available at: [www.nimh.nih.gov/SuicidePrevention/suifact.cfm](http://www.nimh.nih.gov/SuicidePrevention/suifact.cfm). Accessibility verified October 25, 2005.
126. Pepitone-Arreola-Rockwell F, Rockwell D, Core N. Fifty-two medical student suicides. *Am J Psychiatry.* 1981;138:198-201.
127. Hays L, Cheever T, Patel P. Medical student suicide, 1989-1994. *Am J Psychiatry.* 1996;153:553-555.
128. Kessler RC, Borges G, Walters EE. Prevalence of and risk factors for lifetime suicide attempts in the National Comorbidity Survey. *Arch Gen Psychiatry.* 1999;56:617-626.
129. Crosby AE, Cheltenham MP, Sacks JJ. Incidence of suicidal ideation and behavior in the United States, 1994. *Suicide Life Threat Behav.* 1999;29:131-140.
130. Hem E, Gronvold NT, Aasland OG, Ekeberg O. The prevalence of suicidal ideation and suicidal attempts among Norwegian physicians: results from a cross-sectional survey of a nationwide sample. *Eur Psychiatry.* 2000;15:183-189.

131. Tyssen R, Hem E, Vaglum P, Gronvold NT, Ekeberg O. The process of suicidal planning among medical doctors: predictors in a longitudinal Norwegian sample. *J Affect Disord*. 2004;80:191-198.
132. Roy A. Suicide in doctors. *Psychiatr Clin North Am*. 1985;8:377-387.
133. Lepage A, Hunt S. The problem of quality of life in medicine. *JAMA*. 1987;278:47-50.
134. Spilker B, ed. *Quality of Life and Pharmacoeconomics in Clinical Trials*. 2nd ed. New York, NY: Raven Press; 1996.
135. Association of American Medical Colleges. *Physicians for the Twenty-First Century: The GPEP Report: Report of the Panel on the General Professional Education of the Physician and College Preparation for Medicine*. Washington, DC: Association of American Medical Colleges; 1984.
136. Murr AH, Miller C, Papadakis M. Mentorship through advisory colleges. *Acad Med*. 2002;77:1172-1173.
137. Malik S. Students, tutors and relationships: the ingredients of a successful student support scheme. *Med Educ*. 2000;34:635-641.
138. Coles C. Support for medical students in the United Kingdom. *Med Educ*. 1993;27:186-187.
139. Strayhorn G. Effect of a major curriculum revision on students' perceptions of well-being. *Acad Med*. 1989;64:25-29.
140. Murdoch-Eaton DG, Levene MI. Formal appraisal of undergraduate medical students: is it worth the effort? *Med Teach*. 2004;26:28-32.
141. Wolf TM, Randall HM, Faucett JM. A survey of health promotion programs in U.S. and Canadian medical schools. *Am J Health Promot*. 1988;3:33-36.
142. Mouret GM. Stress in a graduate medical degree. *Med J Aust*. 2002;177(suppl):S10-S11.
143. Myers DG. The funds, friends, and faith of happy people. *Am Psychol*. 2000;55:56-67.
144. Lee J, Graham AV. Students' perception of medical school stress and their evaluation of a wellness elective. *Med Educ*. 2001;35:652-659.
145. Rathbun J. Helping medical students develop lifelong strategies to cope with stress. *Acad Med*. 1995;70:955-956.
146. Robins LS, Fantone JC, Oh MS, Alexander GL, Shlafer M, Davis WK. The effect of pass/fail grading and weekly quizzes on first-year students' performances and satisfaction. *Acad Med*. 1995;70:327-329.
147. Dietrick JA, Weaver MT, Merrick HW. Pass/fail grading: a disadvantage for students applying for residency. *Am J Surg*. 1991;162:63-66.
148. Lloyd DA. Pass-fail grading fails to meet the grade. *Acad Med*. 1992;67:583-584.
149. Vosti KL, Jacobs CD. Outcome measurement in postgraduate year one of graduates from a medical school with a pass/fail grading system. *Acad Med*. 1999;74:547-549.
150. Burack JH, Irby DM, Carlone JD, Root RK, Larson EB. Teaching compassion and respect: attending physicians' responses to problematic behaviors. *J Gen Intern Med*. 1999;14:49-55.
151. Freedborn D. Satisfaction, commitment, and psychological well-being among HMO physicians. *Permanente J*. 2000;2:22-30.
152. Pasnau RO, Stoessel P. Mental health service for medical students. *Med Educ*. 1994;28:33-39.
153. Rodolfa E, Chavoor S, Velasquez J. Counseling services at the University of California, Davis: helping medical students cope. *JAMA*. 1995;274:1396-1397.
154. Lerner BA. Students' use of psychiatric services: the Columbia experience. *JAMA*. 1995;274:1398-1399.
155. Keil A. AIMS Committees: a resource for medical students. *JAMA*. 1995;274:1399.
156. Hays LR, Dickson LR, Lyles MR, Ludwig AM, Martin CA, Bird MA. Treating psychiatric problems in medical students. *Am J Psychiatry*. 1986;143:1428-1431.
157. Ackerman TF, Wall HP. A programme for treating chemically dependent medical students. *Med Educ*. 1994;28:40-46.
158. Kelly JA, Bradlyn AS, Dubbert PM, St Lawrence JS. Stress management training in medical school. *J Med Educ*. 1982;57:91-99.
159. Holtzworth-Munroe A, Munroe MS, Smith RE. Effects of a stress-management training program on first- and second-year medical students. *J Med Educ*. 1985;60:417-419.
160. McCue JD, Sachs CL. A stress management workshop improves residents' coping skills. *Arch Intern Med*. 1991;151:2273-2277.
161. Branch W, Pels RJ, Lawrence RS, Arky R. Becoming a doctor: critical-incident reports from third-year medical students. *N Engl J Med*. 1993;329:1130-1132.
162. Pololi L, Frankel RM. Small-group teaching emphasizing reflection can positively influence medical students' values. *Acad Med*. 2001;76:1172-1173.
163. Gaber RR, Martin DM. Still-Well osteopathic medical student wellness program. *J Am Osteopath Assoc*. 2002;102:289-292.
164. Ware JE, Kosinski M, Dewey JE, Gandek B. *How to Score and Interpret Single-Item Health Status Measures: A Manual for Users of the SF-8 Health Survey*. Lincoln, RI: QualityMetric Incorporated, 2001.
165. Davis M. A multidimensional approach to individual differences in empathy. *JSAS Catalog of Selected Documents in Psychology*. 1980;10, 85.
166. Shanafelt TD, West C, Zhao X, et al. Relationship between increased personal well-being and enhanced empathy among internal medicine residents. *J Gen Intern Med*. 2005;20:559-564.
167. Sloan JA, Loprinzi CL, Kuross SA, et al. Randomized comparison of four tools measuring overall quality of life in patients with advanced cancer. *J Clin Oncol*. 1998;16:3662-3673.
168. Grunberg SM, Groshen S, Steingass S, Zaretsky S, Meyerowitz B. Comparison of conditional quality of life terminology and visual analogue scale measurements. *Qual Life Res*. 1996;5:65-72.
169. Gudex C, Dolan P, Kind P, Williams A. Health state valuations from the general public using the visual analogue scale. *Qual Life Res*. 1996;5:521-531.
170. Hyland ME, Sodergren SC. Development of a new type of global quality of life scale, and comparison of performance and preference for 12 global scales. *Qual Life Res*. 1996;5:469-480.
171. Gill P, Kaur JS, Rummans T, Novotny PJ, Sloan JA. The hospice patient's primary caregiver: what is their quality of life? *J Psychosom Res*. 2003;55:445-451.